



The genomic architecture of segmental duplications and associated copy number variants in dogs

Thomas J. Nicholas, Ze Cheng, Mario Ventura, et al.

Genome Res. published online January 7, 2009

Access the most recent version at doi:[10.1101/gr.084715.108](https://doi.org/10.1101/gr.084715.108)

| | |
|-------------------------------|---|
| P<P | Published online January 7, 2009 in advance of the print journal. |
| Accepted Preprint | Peer-reviewed and accepted for publication but not copyedited or typeset; preprint is likely to differ from the final, published version. |
| Email alerting service | Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here |

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *Genome Research* go to:
<http://genome.cshlp.org/subscriptions>

The genomic architecture of segmental duplications and associated copy number variants in dogs

Thomas J. Nicholas¹, Ze Cheng¹, Mario Ventura², Katrina Mealey³, Evan E. Eichler^{1,4*}, and Joshua M. Akey^{1*}

¹ Department of Genome Sciences, University of Washington, 1705 NE Pacific, Seattle, WA. 98195, USA.

² Department of Genetics and Microbiology, University of Bari, Via Amendola 165/A, 70124, Bari, Italy

³ Department of Veterinary Clinical Sciences, College of Veterinary Medicine, Washington State University, Pullman WA 99164-6610, USA.

⁴ Howard Hughes Medical Institute, Seattle, Washington, USA.

Running Title: Structural variation in dogs

To whom correspondence should be addressed:

Joshua M. Akey
Department of Genome Sciences
University of Washington
Foege Building S-303
1705 NE Pacific St
Seattle WA 98195-5065
Phone: (206) 543-7254
Fax: (206) 685-7301
e-mail: akeyj@u.washington.edu

or

Evan E. Eichler
Department of Genome Sciences
University of Washington
Foege Building S-413
1705 NE Pacific St
Seattle WA 98195-5065
Phone: (206) 543-9526
Fax: (206) 685-7301
e-mail: eee@u.washington.edu

Abstract

Structural variation is as an important and abundant source of genetic and phenotypic variation. Here we describe the first systematic and genome-wide analysis of segmental duplications and associated copy number variants (CNVs) in the modern domesticated dog, *Canis familiaris*, which exhibits considerable morphological, physiological, and behavioral variation. Through computational analyses of the publicly available canine reference sequence, we estimate that segmental duplications comprise approximately 4.21% of the canine genome. Segmental duplications overlap 841 genes, and are significantly enriched for specific biological functions such as immunity and defense and KRAB-box transcription factors. We designed high-density tiling arrays spanning all predicted segmental duplications and performed aCGH in a panel of 17 breeds and a gray wolf. In total, we identified 3,583 CNVs, approximately 68% of which were found in two or more samples, that map to 678 unique regions. CNVs span 429 genes that are involved in a wide variety of biological processes such as olfaction, immunity, and gene regulation. Our results provide insight into mechanisms of canine genome evolution and generate a valuable resource for future evolutionary and phenotypic studies.

Introduction

The unique evolutionary history of domesticated dogs (*Canis familiaris*), including strong artificial selection, population bottlenecks, and inbreeding, has resulted in over 400 genetically distinct breeds that make them well suited for addressing fundamental questions in population genetics, evolution, and the genetic architecture of phenotypic variation. In particular, domesticated dogs have engendered considerable interest because, since their domestication over 14,000 years ago (Vila et al. 1997; Leonard et al. 2002; Savolainen et al. 2002), they have become one of the most phenotypically diverse mammalian species with an incredible assortment of shapes, sizes, and temperaments (Neff and Rine 2006). Beyond curiosity in outward appearances, canine genetics is also relevant to human health as dogs are afflicted with over 350 inherited diseases (Patterson et al. 1988), many of which are similar to human diseases.

A number of enabling resources for canine genomics have recently become available, including the development of an integrated canine linkage-radiation hybrid map (Mellersh et al. 2000), a 7.5X high-quality reference genome sequence (Lindblad-Toh et al. 2005), the construction of a dense map of over 2.5 million SNPs identified in a diverse panel of breeds (Lindblad-Toh et al. 2005), and the development of SNP genotyping arrays (Karlsson et al. 2007). These resources have provided important foundations for delimiting patterns of population structure among breeds (Irion et al. 2003; Parker et al. 2004; Karlsson et al. 2007; Quignon et al. 2007), inferring targets of artificial selection (Pollinger et al. 2005), and mapping traits such as Collie Eye Anomaly (Parker et al. 2007), body size (Sutter et al. 2007), and muscle mass (Mosher et al. 2007).

In contrast to SNPs and microsatellites, structural variation has received considerably less attention in dogs. Changes in DNA content are a significant source of genetic and phenotypic variation between individuals (Emanuel and Shaikh 2001; Bailey and Eichler

2006; Feuk, Carson, and Scherer 2006; Beckmann, Estivill, and Antonarakis 2007; Conrad and Antonarakis 2007; Sebat 2007). Segmental duplications in particular are substrates of genome innovation, genomic rearrangements, and hotspots of CNV formation (Sharp et al. 2005; Graubert et al. 2007; She et al. 2008). Although segmental duplications and CNVs have been extensively studied in other organisms (Bailey et al. 2001; Bailey et al. 2002; Bailey et al. 2004; Iafrate et al. 2004; Tuzun, Bailey, and Eichler 2004; Cheng et al. 2005; Sharp et al. 2005; Tuzun et al. 2005; Conrad et al. 2006; Goidts et al. 2006; McCarroll et al. 2006; Perry et al. 2006; Redon et al. 2006; Graubert et al. 2007; Guryev et al. 2008; Perry et al. 2008; She et al. 2008), to date no such analyses have been performed in dogs. Recent studies demonstrate the potential contribution of CNVs to specific canine morphological phenotypes, such as dorsal hair ridge in Rhodesian and Thai Ridgebacks (Salmon Hillbertz et al. 2007). Thus, a more comprehensive understanding of the full spectrum of canine genomic variation is important for unraveling the genetic basis of variation in morphological, physiological, behavioral, and disease phenotypes segregating within and between breeds (Neff and Rine 2006).

Here we describe the first genome-wide and systematic analysis of segmental duplications and their associated CNVs in dogs. We find that similar to other mammalian genomes, recent segmental duplications comprise an appreciable fraction of the canine genome. Using high-density aCGH experiments specifically designed to interrogate putative segmental duplications, we identified 3,583 CNVs in a panel of 17 genetically and phenotypically diverse breeds and a gray wolf.

Results and Discussion

Genome-wide Identification and Organization of Segmental Duplications

We applied two well-established computational approaches, Whole Genome Shotgun Sequence Detection (WSSD) (Bailey et al. 2002) and Whole Genome Assembly Comparison (WGAC) (Bailey et al. 2001), to the publicly available canine genome sequence assembly (CanFam2.0) to detect putative segmental duplications. Briefly, WGAC identifies paralogous sequences ≥ 1 kb in length with $\geq 90\%$ sequence identity and WSSD identifies genomic regions that exhibit significant depth of coverage by aligning whole genome shotgun sequencing reads to the reference genome sequence (see Methods). Using these computational algorithms, we predict 9,137 segmental duplications, spanning approximately 106.6 Mb of DNA sequence (Figure 1; Supplementary Table 1). The average size of predicted segmental duplications is approximately 11.7 kb (sd = 24.9 kb). We estimate that recent segmental duplications comprise approximately 4.21% of the canine reference genome, which is consistent with similar observations in human and mouse (Bailey et al. 2001; Bailey et al. 2002; Bailey et al. 2004; She et al. 2008). As expected, the “uncharacterized chromosome” (chrUn), which consists of sequence that cannot be uniquely mapped to the genome, contains the majority of predicted duplication bases (65%).

Furthermore, similar to humans and mice, there is a greater proportion of intrachromosomal versus interchromosomal duplications, with approximately 60% of predicted duplications being intrachromosomal. Pericentromeric regions represent 3.4% of genomic sequence, but show an enrichment of 3-fold for duplications (p-value < 0.001) and contain 10.3% of all duplicated bases. Similarly, subtelomeric regions show an enrichment of 2.3-fold (p-value < 0.001) and contain 7.9% of duplicated bases.

841 genes were located in predicted segmental duplications. In order to test the hypothesis that particular gene classes are overrepresented in duplicated regions, we assigned

PANTHER Molecular Function terms to all genes that overlapped duplications. Statistically significant enrichment was observed for seven categories (Supplementary Table 2).

Consistent with similar analyses of duplications in other organisms (Bailey et al. 2002; Bailey et al. 2004; Tuzun, Bailey, and Eichler 2004; Sharp et al. 2005), we observe significant enrichment in genes that participate in defense/immunity, receptors, and signaling (Supplementary Table 2). Interestingly, KRAB box transcription factors were also significantly enriched among our set of predicted duplications (Supplementary Table 2), which has not been previously observed in duplications from additional species. KRAB box transcriptional factors are part of a large gene family that are believed to bind to DNA and exhibit transcriptional repression (Urrutia 2003). While the precise phenotypes that KRAB box transcription factors contribute to are largely unknown, recent work has shown that a particular KRAB box transcription factor influences mouse embryonic morphological developmental (Garcia-Garcia, Shibata, and Anderson 2008). No additional classes of transcription factors showed enrichment.

FISH Characterization of Predicted Segmental Duplication

We experimentally validated a subset of duplicated regions by fluorescent *in situ* hybridization (FISH). 42 large-insert dog BAC clones corresponding to WGAC and WSSD duplicated regions (>20 kb in length) were used as probes and hybridized against a fibroblast *Canis familiaris* cell line (Supplemental Table 3). We observed multiple signals either by examination of interphase or metaphase FISH for 20/42 of probes, including 14 intrachromosomal, 5 interchromosomal, and a single probe that mapped to multiple centromeric regions. Only 1 of the interchromosomal probes showed more than 3 distinct signals, while the majority (10/14) of intrachromosomal duplication signals were clustered. Similar to the mouse genome (She et al. 2008), these data suggest that tandem

intrachromosomal duplications predominate in the dog genome (Figure 2). The basis for the remaining 22 BAC probes consistent with single copy sequence is unknown. We note, however, that the breed origin for the *Canis familiaris* cell line used in the FISH experiments is not known, and structural polymorphism as well as limitations of BAC-FISH to detect duplications < 40 kb (especially in the case of tandem duplications) may account for differences between the computational predictions and experimental data.

CNVs and Segmental Duplications

We designed a custom high-density tiling array covering all regions with significant WGAC or WSSD support to identify CNVs. Note, this includes both the 106.6 Mb of sequence predicted to be segmental duplications (both WGAC and WSSD support) as well as an additional 16.4 Mb of sequence with either WGAC or WSSD support. Obviously, our study design will not detect CNVs located outside of these regions, but we focused on them for two reasons. First, previous studies have demonstrated that segmental duplications are enriched 4-20 fold for CNVs (Iafrate et al. 2004; Sebat et al. 2004; Sharp et al. 2005; Tuzun et al. 2005; Perry et al. 2006; Graubert et al. 2007; Guryev et al. 2008; She et al. 2008). Second, by restricting our aCGH experiments to regions most likely to harbor CNVs we were able to dramatically increase probe density (average probe spacing of 200 bp) resulting in a more exhaustive discovery of smaller CNVs as well as increasing the robustness of individual CNV calls.

The remaining probes were placed in 5 kb flanking regions and putative single copy control regions with a mean spacing of 1 kb and 350 bp respectively. Thus, in total over 137 Mb of sequence was studied. In total, 19 aCGH experiments were performed, each using a common reference sample derived from a female boxer. Specifically, hybridizations were performed on 17 diverse breeds (see Methods), a gray wolf, and a self-self hybridization.

We used a previously described hidden markov model method to identify changes in \log_2 signal intensity corresponding to gains and losses in copy number (Rueda and Diaz-Uriarte 2007). Using conservative criteria (see Methods), 3,583 CNVs were identified (1,578 gains and 2,005 losses) that map to 678 distinct regions (Figure 3; Supplementary Table 4). CNVs comprise 24 Mb of polymorphic sequence and approximately 20% of the predicted segmental duplications exhibit CNVs (Figure 1). Interestingly, approximately 50% of CNVs exhibit both WGAC and WSSD support.

The average number of CNVs per breed was 199, ranging from 118 (German Shorthaired Pointer) to 298 (Basenji), and approximately 68% of CNVs were found in two or more individuals (Table 1). In contrast, only one CNV was called in the self-self hybridization and no CNVs were called in the single copy control regions (Figure 3). We also identified CNVs using an additional algorithm and found good overlap (Supplementary Note 1). Thus, these observations suggest that the false discovery rate (FDR) among the set of predicted CNVs is low (~3%), although technical issues such as sequence divergence of individual dogs relative to the reference genome sequence or heterogeneity in DNA quality among samples makes it difficult to precisely quantify the FDR.

In general, the number of CNVs identified in each sample is consistent with previous estimates (where available) of breed specific founding and effective population sizes (Calboli et al. 2008) and levels of polymorphism based on 27,000 SNPs (Karlsson et al. 2007). Of interest, 169 CNVs were identified in the Boxer (Tasha) whose DNA was selected for the dog genome project (Lindblad-Toh et al. 2005) based on its relatively low level of genetic diversity. The somewhat lower level of polymorphism in this boxer, however, does not necessarily imply genetic homogeneity within boxers or other breeds (Parker et al. 2004; Sutter and Ostrander 2004; Wayne and Ostrander 2007). Indeed, a number of recent data sets and analyses suggest considerable genetic diversity exists within breeds (Quignon et al. 2007;

Bjornerfeldt et al. 2008), which is consistent with our observation of segregating CNVs among boxers.

qPCR Analysis of Two CNV Regions

Quantitative PCR (qPCR) was performed using Taqman probes and primers on all dogs used in the aCGH experiments to further validate CNV regions and individual calls (see Methods). Probes were targeted to two regions that show homology to the human *GCKR* and *PHYH* genes. As shown in Figure 4, qPCR identified 16 and 15 individuals with copy number differences relative to the reference sample for *GCKR* and *PHYH*, respectively. Assuming the qPCR results represent the true copy number status of individual dogs, a total of two false positives occurred in the aCGH data (zero for *GCKR* and two for *PHYH*). Thus, the qPCR data confirms the designation of *GCKR* and *PHYH* as CNV regions, and suggests a low FDR of actual calls within such regions ($2/31 = 6.5\%$). The qPCR data also confirm the conservativeness of thresholds for calling CNVs in the aCGH data, as 11 individuals were identified with copy number changes in qPCR but not in the aCGH data (false negative rate $\sim 30\%$).

Fine-scale Architectural Complexity of CNVs

Correctly delineating CNV boundaries is important for both understanding the molecular mechanisms governing CNV formation and correlating copy number changes with phenotypic variation (Perry et al. 2008). The high probe density of our tiling arrays allowed us to investigate patterns of breakpoint variation across individuals. Of the 460 distinct CNV regions where a gain or loss was observed in two or more samples, 235 exhibited relatively simple architectures with consistent patterns of breakpoints across individuals.

The remaining 225 CNV regions (~50%) showed fine-scale architectural complexity in the form of substantial inter-individual variation in breakpoints or spatial heterogeneity in copy number. Similarly complex CNV regions have been described (Redon et al. 2006; Perry et al. 2008; She et al. 2008), primarily in studies focusing on segmental duplications (Goidts et al. 2006). A particularly interesting pattern was observed for 20 CNV regions where alternating gains and losses occurred within individual dogs. An example of one such region spanning over 400 kb on chromosome 17 is shown in Figure 5. In this region, we observed dogs that have gains or losses across the whole region, both gains and losses, and no copy number change relative to the reference sequence (Figure 5). The precise mechanistic basis for such complex CNV patterns is unclear, and may be attributable to non-allelic homologous recombination or less understood mechanisms such as the recently proposed replication based fork stalling and template switching model (Lee, Carvalho, and Lupski 2007).

Gene Content of CNV Regions

CNVs overlap 429 genes (Supplementary Table 5), 415 of which span the complete coding region. The set of copy number variable genes possess a wide spectrum of PANTHER molecular functions (Supplementary Tables 2 and 5), and provides a rich resource for testing hypotheses on the genetic basis of phenotypic variation within and among breeds. For example, in humans, copy number variation of cytochrome P450 genes, such as *CYP2D6*, contributes to inter-individual variation in drug metabolism phenotypes (Daly 2004; Ledesma and Agundez 2005; Ouahchi, Lindeman, and Lee 2006). Similar to humans, adverse drug responses have been described in dogs, which often show marked variation in prevalence between breeds (Hickford, Barr, and Erb 2001; Mealey et al. 2001; Mealey, Northrup, and Bentjen 2003; Nelson et al. 2003; Neff et al. 2004; Trepanier 2004). Several CYP genes overlap CNVs, perhaps the most interesting of which is *CYP3A12* the canine ortholog to

human *CYP3A4*, which is the most abundant hepatic and intestinal cytochrome P450 isoform and is involved in metabolizing a substantial fraction of all drugs (Schuetz 2004).

Specifically, nine dogs (Afghan Hound, Doberman Pinscher, German Shepherd, Labrador Retriever, Rottweiler, West Highland White Terrier, Yorkshire Terrier, Boxer and Wolf) show partial loss of *CYP3A12*, and of these adverse drug responses have been described in Doberman Pinschers (sulfonamide hypersensitivity; Trepanier 2004), Labrador Retrievers (carprofen-induced hepatic toxicity; Hickford, Barr, and Erb 2001), and Boxers (acepromazine sensitivity, although this result is controversial; Wagner, Wright, and Hellyer 2003). Obviously, these observations, while interesting, require additional study to better delimit the relationship between *CYP3A12* copy number and variation in drug metabolism phenotypes.

CNVs that span potential genes influencing disease susceptibility were also identified. For instance, the glucokinase regulatory protein gene (*GCKR*) is located in a complex CNV region (Figure 5). Recent genome-wide association studies in humans have found that *GCKR* variation increases susceptibility to type 2 diabetes (Saxena et al. 2007). In our panel of dogs, there is a suggestive pattern of *GCKR* copy number status and risk of developing diabetes mellitus that warrants further study, with breeds considered at high risk preferentially showing deletions for varying parts of *GCKR* (Supplementary Table 6).

Conclusions

In summary, we have described the first genome-wide analysis of segmental duplications and associated CNVs in the modern domesticated dog. We found extensive copy number variation in segmental duplications across 17 phenotypically diverse breeds that affect 429 genes. Our study provides the foundation for correlating structural variation with phenotypic variation observed within and between breeds, which we suspect will be an

important complement to SNP centric genome-wide association studies (Karlsson et al. 2007; Mosher et al. 2007; Parker et al. 2007; Quignon et al. 2007; Salmon Hillbertz et al. 2007; Sutter et al. 2007). However, in order to perform more principled phenotypic studies with structural variation, it will be necessary to better delimit the population genetic characteristics of CNVs including a broader sampling of breeds, extending CNV identification and characterization to regions of the genome outside of segmental duplications, assessing allele frequency distributions in unrelated individuals within breeds, and determining levels of linkage disequilibrium between CNVs and SNPs. Ultimately, understanding how the full spectrum of canine genomic variation influences phenotypic variation will provide insight into the genetic architecture of phenotypes and mechanisms of rapid short-term evolutionary change.

Methods

Computational analysis of segmental duplications

We downloaded CanFam2.0 genomic sequence data from the UCSC Genome Browser (<http://genome.ucsc.edu/>) and whole genome shotgun sequence (WGS) reads were obtained through NCBI (<http://www.ncbi.nlm.nih.gov/>). WGAC and WSSD were performed as previously described (Bailey et al. 2001; Bailey et al. 2002). Briefly, WGAC identifies paralogous stretches of sequence by fragmenting the genome into 400 kb fragments, which are masked for repeats with RepeatMasker and removed. Global alignments of these fragments are performed with alignments of >90% sequence identity and >1 kb in length deemed as paralogs. Masked regions are then reinserted and fragments are returned to their genomic locations.

WSSD identifies genomic regions of significant depth of coverage by aligning WGS reads to the reference genome sequence. We initially conducted WSSD on a set of 13 training BACs (also obtained from NCBI; AC090032.2, AC090890.3, AC091119.2, AC117937.4, AC147678.5, AC147681.8, AC147784.3, AC090889.3, AC090972.5, AC092249.3, AC147677.4, AC147680.4, AC147707.5). BACs were masked for repeats and MegaBLAST alignments of these BACs were performed against a database of WGS reads. We calculated duplication depth by counting the number of WGS reads aligning to 5 kb sliding windows. In addition, we calculated nucleotide divergence between the WGS reads and the BAC sequences for each 5 kb window. The distribution of alignment depth and divergence in this training set allows empirical thresholds to be determined. Consistent with previous studies (Bailey et al. 2002; Bailey et al. 2004; Cheng et al. 2005; She et al. 2008), we define significant alignment depth and divergence scores as those that are greater than 3 standard deviations from the mean. After training, we masked the entire canine reference genome for common repeats with <10% divergence and performed MegaBLAST alignments of the WGS

reads to the reference genome. Following previous studies (She et al. 2008), we defined segmental duplications based on the union of significant WGAC hits with less than 94% sequence identity and WSSD results (see Supplementary Table 1).

Bioinformatics analysis of segmental duplications

We investigated the genomic distribution of segmental duplications by testing the hypothesis that pericentromeric and subtelomeric regions were enriched for duplications (Bailey et al. 2001). Since the pericentromeric and subtelomeric regions are not well annotated we defined pericentromeric and subtelomeric regions as 2 Mb from the most centromeric base and 2 Mb from the end(s) of chromosomes, respectively. Since all dog chromosomes are acrocentric, with the exception of the X chromosome, this results in a 2 Mb pericentromeric region at one end of the chromosome and a 2 Mb subtelomeric region at the other end of the chromosome. In the case of the X chromosome, the pericentromeric region was defined as two 1 Mb regions that flank the centromeric region and two 1 Mb subtelomeric ends on both ends of the chromosome. No sequence from chrUn was included. All predicted duplicated bases that overlap these regions were totaled and chi-square tests were used to test the null hypothesis of no enrichment as previously described (Bailey et al. 2001).

We obtained a catalog of all canine peptides from Ensembl (ftp://ftp.ensembl.org/pub/current_fasta/canis_familiaris/pep/). This yielded 25,546 peptides, 1078 of which overlap with predicted segmental duplications, and correspond to 841 unique Ensembl genes. PANTHER Molecular Function terms were assigned to all peptides using the PANTHER Hidden Markov Model scoring tools (<http://www.pantherdb.org/downloads/>). PANTHER Molecular Function terms with less than five observations among the duplicated genes were not analyzed further. Similar analyses were performed on the 547 peptides

(corresponding to 429 Ensembl genes) that overlap with the 678 unique CNV regions. We tested the hypothesis that the remaining PANTHER Molecular Function terms were overrepresented in segmental duplications and CNVs with the hypergeometric distribution. Bonferroni corrections were used to correct p-values for multiple hypothesis testing.

DNA samples

Breeds used in this study include an Afghan Hound, Basenji, Belgian Shepherd (Tervuren), Boxer, Doberman Pinscher, German Shepherd, German Shorthaired Pointer, Golden Retriever, Labrador Retriever, Pug, Rottweiler, Shetland Sheepdog, Siberian Husky, Standard Poodle, West Highland White Terrier, Whippet, and Yorkshire Terrier. In addition, genomic DNA was obtained from Tasha, a boxer whose DNA was used in the canine genome project, and also a gray wolf. Sample collection was approved by the Animal Care and Use Committees from Washington State University. We isolated genomic DNA from whole blood samples using Qiagen's QIAamp DNA Blood Maxi Kit. All dogs were screened for correct AKC status. We assessed DNA quality and purity by OD 260/280 and OD 260/230 readings and by digesting genomic DNA with a salt sensitive restriction enzyme (NlaIII).

FISH and image analysis

Canis familiaris metaphase spreads were prepared from PHA-stimulated peripheral lymphocytes of a normal donor of unknown breed identity by standard procedures. DNA extraction from BACs has already been reported. FISH experiments were performed essentially as previously described (Ventura et al. 2003). Briefly, DNA probes were directly labeled with Cy3-dUTP (Perkin-Elmer) by nick-translation. Two hundred nanograms of labeled probe was used for the FISH experiments. Hybridization was performed at 37°C in 2 X SSC, 50% (v/v) formamide, 10% (w/v) dextran sulfate and 3 mg of sonicated salmon sperm DNA, in a volume of 10 µL. Post-hybridization washing was at 60°C in 0.1XSSC

(three times, high stringency).

Digital images were obtained using a Leica DMRXA epifluorescence microscope equipped with a cooled CCD camera (Princeton Instruments). Cy3 (red) and DAPI (blue) fluorescence signals, detected with specific filters, were recorded separately as grayscale images. Pseudocoloring and merging of images were performed using Adobe Photoshop software.

aCGH chip design and hybridizations

The production of the aCGH chip was carried out by Nimblegen Systems Inc (www.NimbleGen.com) with final probe design and locations being approved after individual inspection by eye. We targeted probes to regions designated as segmental duplications (significant WGAC and WSSD support; 106.6 Mb), as well as all regions not formally designated as segmental duplications but possessing significant WGAC or WSSD support (16.4 Mb). In total, 368,360 probes were placed into these regions (spanning 123 Mb) with an average probe spacing of 200 bp. Each segmental duplication region was flanked with 5kb of putative single copy sequence. In cases where the added flanks resulted in overlaps between separate segmental duplications, the segmental duplications were merged together with new flanking regions added to the merged segmental duplication. A total of 9,278 probes were placed in flanking regions (10.6 Mb) with a mean spacing of 1 kb. Finally, we placed 8,790 probes with a mean spacing of 350 bp into single copy control regions (3.5 Mb). Single copy control regions were defined as being at least 5 Mb away from any predicted segmental duplications (with the exception of 3 control regions on the X chromosome which were at least 2.5 Mb away from any predicted segmental duplication). All genomic DNA samples were sent to Nimblegen for the hybridizations to be performed. A female Boxer distinct from Tasha was used as a reference sample in each hybridization.

aCGH data analysis and CNV calling

For each hybridization, normalized \log_2 ratios were first averaged across 2 kb windows. Due to the unique assembly of chrUn sequences, each chrUn assembly contig was analyzed as if it were an individual chromosome and separately from the assembled chromosomes. All averaged 2 kb windows were analyzed using the R package RJaCGH (Rueda and Diaz-Uriarte 2007). We used the self-self hybridization and control regions to define suitable thresholds to apply to the RJaCGH calls in order to minimize false positives. Specifically, we retained predicted CNVs if it had at least 5 datapoints supporting it, thus limiting the minimum CNV size to approximately 10 kb. The chrUn results were further filtered by requiring the average \log_2 value of the CNV to be greater than 0.25 or less than -0.25. We merged overlapping CNV coordinates across hybridizations to form unique CNV regions. All unique CNV regions on the X chromosome that were only supported by male dogs were removed to avoid potential complications since the reference sample was a female. We repeated the entire RJaCGH analyses for a subset of the arrays to ensure consistency in calls. We assessed breakpoint variation by analyzing each individual CNV in a defined unique region and determining whether its length varied by more than 5 kb from the entire CNV region length.

The false discovery rate (FDR) was estimated based on the observation of a single false positive in the self-self hybridization and thus a rough estimate of the FDR is the expected number of false positives per array (1) times the number of total arrays (18) divided by the total number of unique CNV regions (678), resulting in an estimated FDR of 2.65%. Note, this calculation is only approximate because it assumes that each false positive results in a unique CNV region and does not take into account the potential for varying false positive rates across arrays.

qPCR

We performed qPCR on all dogs from two CNV regions (*GCKR* and *PHYH*). Specifically, Taqman probes and primers (Applied Biosystems) were designed for three regions; a single copy control region (labeled with FAM dye) and two test regions (each labeled with VIC dye). Primer and probe sequences are available upon request. Assays were performed on an ABI 7900HT (Applied Biosystems) using 20- μ l reactions containing 10- μ l of Taqman Universal PCR Master Mix, 250 nM of FAM probe, 900 nM of forward and reverse primers for FAM probe, 250 nM of VIC probe, 900 nM of forward and reverse primers for VIC probe, and 30 ng of genomic DNA. Amplification was done under the following conditions: one cycle at 50°C for 2 min, one cycle at 95°C for 10 min, 40 cycles at 95°C for 15s and 60°C for 1 min. Serial dilutions were performed for each assay to estimate the PCR efficiency (E). Using the ΔC_T method, relative copy number was determined with respect to the same reference Boxer sample used in the aCGH experiments. C_T values were adjusted for PCR efficiency (E) as $\log_2(E^{C_T})$. The C_T values were then normalized by subtracting the VIC C_T value from the FAM C_T value (FAM C_T – VIC C_T). The relative copy number was determined as $2^{-(\text{normalized } C_T \text{ for test strain} - \text{normalized } C_T \text{ for reference strain})}$. In total, 6 independent replicates were performed for each individual. Statistical significance was determined by a one sample t-test. Similar results were obtained with a one sample Wilcoxon test (data not shown).

Data Release

All aCGH data has been submitted to the gene expression omnibus (<http://www.ncbi.nlm.nih.gov/geo/>) under the accession number GSE13266.

Acknowledgements

We thank pet owners, and their dogs, for participating in this research, Dayna Akey for helpful discussions, and Kerstin Lindblad-Toh for providing DNA for the sample Boxer (Tasha) used to construct the reference dog genome sequence. This work was supported in part by grants from the University of Washington Royalty Research Fund (JMA), the American Kennel Club (JMA and KM), Howard Hughes Medical Institute (EEE), and a Sloan Fellowship in Computational Biology (JMA).

References

- Bailey, J.A., D.M. Church, M. Ventura, M. Rocchi, and E.E. Eichler. 2004. Analysis of segmental duplications and genome assembly in the mouse. *Genome Res* **14**: 789-801.
- Bailey, J.A. and E.E. Eichler. 2006. Primate segmental duplications: crucibles of evolution, diversity and disease. *Nat Rev Genet* **7**: 552-564.
- Bailey, J.A., Z. Gu, R.A. Clark, K. Reinert, R.V. Samonte, S. Schwartz, M.D. Adams, E.W. Myers, P.W. Li, and E.E. Eichler. 2002. Recent segmental duplications in the human genome. *Science* **297**: 1003-1007.
- Bailey, J.A., A.M. Yavor, H.F. Massa, B.J. Trask, and E.E. Eichler. 2001. Segmental duplications: organization and impact within the current human genome project assembly. *Genome Res* **11**: 1005-1017.
- Beckmann, J.S., X. Estivill, and S.E. Antonarakis. 2007. Copy number variants and genetic traits: closer to the resolution of phenotypic to genotypic variability. *Nat Rev Genet* **8**: 639-646.
- Bjornerfeldt, S., F. Hailer, M. Nord, and C. Vila. 2008. Assortative mating and fragmentation within dog breeds. *BMC Evol Biol* **8**: 28.
- Calboli, F.C., J. Sampson, N. Fretwell, and D.J. Balding. 2008. Population structure and inbreeding from pedigree analysis of purebred dogs. *Genetics* **179**: 593-601.
- Cheng, Z., M. Ventura, X. She, P. Khaitovich, T. Graves, K. Osoegawa, D. Church, P. DeJong, R.K. Wilson, S. Paabo et al. 2005. A genome-wide comparison of recent chimpanzee and human segmental duplications. *Nature* **437**: 88-93.
- Conrad, B. and S.E. Antonarakis. 2007. Gene duplication: a drive for phenotypic diversity and cause of human disease. *Annu Rev Genomics Hum Genet* **8**: 17-35.
- Conrad, D.F., T.D. Andrews, N.P. Carter, M.E. Hurles, and J.K. Pritchard. 2006. A high-resolution survey of deletion polymorphism in the human genome. *Nat Genet* **38**: 75-81.
- Daly, A.K. 2004. Pharmacogenetics of the cytochromes P450. *Curr Top Med Chem* **4**: 1733-1744.
- Emanuel, B.S. and T.H. Shaikh. 2001. Segmental duplications: an 'expanding' role in genomic instability and disease. *Nat Rev Genet* **2**: 791-800.
- Feuk, L., A.R. Carson, and S.W. Scherer. 2006. Structural variation in the human genome. *Nat Rev Genet* **7**: 85-97.
- Garcia-Garcia, M.J., M. Shibata, and K.V. Anderson. 2008. Chato, a KRAB zinc-finger protein, regulates convergent extension in the mouse embryo. *Development* **135**: 3053-3062.
- Goidts, V., D.N. Cooper, L. Armengol, W. Schempp, J. Conroy, X. Estivill, N. Nowak, H. Hameister, and H. Kehrer-Sawatzki. 2006. Complex patterns of copy number variation at sites of segmental duplications: an important category of structural variation in the human genome. *Hum Genet* **120**: 270-284.
- Graubert, T.A., P. Cahan, D. Edwin, R.R. Selzer, T.A. Richmond, P.S. Eis, W.D. Shannon, X. Li, H.L. McLeod, J.M. Cheverud et al. 2007. A high-resolution map of segmental DNA copy number variation in the mouse genome. *PLoS Genet* **3**: e3.
- Guryev, V., K. Saar, T. Adamovic, M. Verheul, S.A. van Heesch, S. Cook, M. Pravenec, T. Aitman, H. Jacob, J.D. Shull et al. 2008. Distribution and functional impact of DNA copy number variation in the rat. *Nat Genet* **40**: 538-545.
- Hickford, F.H., S.C. Barr, and H.N. Erb. 2001. Effect of carprofen on hemostatic variables in dogs. *Am J Vet Res* **62**: 1642-1646.

- Iafrate, A.J., L. Feuk, M.N. Rivera, M.L. Listewnik, P.K. Donahoe, Y. Qi, S.W. Scherer, and C. Lee. 2004. Detection of large-scale variation in the human genome. *Nat Genet* **36**: 949-951.
- Irion, D.N., A.L. Schaffer, T.R. Famula, M.L. Eggleston, S.S. Hughes, and N.C. Pedersen. 2003. Analysis of genetic variation in 28 dog breed populations with 100 microsatellite markers. *J Hered* **94**: 81-87.
- Karlsson, E.K., I. Baranowska, C.M. Wade, N.H. Salmon Hillbertz, M.C. Zody, N. Anderson, T.M. Biagi, N. Patterson, G.R. Pielberg, E.J. Kulbokas, 3rd et al. 2007. Efficient mapping of mendelian traits in dogs through genome-wide association. *Nat Genet* **39**: 1321-1328.
- Ledesma, M.C. and J.A. Agundez. 2005. Identification of subtypes of CYP2D gene rearrangements among carriers of CYP2D6 gene deletion and duplication. *Clin Chem* **51**: 939-943.
- Lee, J.A., C.M. Carvalho, and J.R. Lupski. 2007. A DNA replication mechanism for generating nonrecurrent rearrangements associated with genomic disorders. *Cell* **131**: 1235-1247.
- Leonard, J.A., R.K. Wayne, J. Wheeler, R. Valadez, S. Guillen, and C. Vila. 2002. Ancient DNA evidence for Old World origin of New World dogs. *Science* **298**: 1613-1616.
- Lindblad-Toh, K. C.M. Wade T.S. Mikkelsen E.K. Karlsson D.B. Jaffe M. Kamal M. Clamp J.L. Chang E.J. Kulbokas, 3rd M.C. Zody et al. 2005. Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature* **438**: 803-819.
- McCarroll, S.A., T.N. Hadnott, G.H. Perry, P.C. Sabeti, M.C. Zody, J.C. Barrett, S. Dallaire, S.B. Gabriel, C. Lee, M.J. Daly et al. 2006. Common deletion polymorphisms in the human genome. *Nat Genet* **38**: 86-92.
- Mealey, K.L., S.A. Bentjen, J.M. Gay, and G.H. Cantor. 2001. Ivermectin sensitivity in collies is associated with a deletion mutation of the *mdr1* gene. *Pharmacogenetics* **11**: 727-733.
- Mealey, K.L., N.C. Northrup, and S.A. Bentjen. 2003. Increased toxicity of P-glycoprotein-substrate chemotherapeutic agents in a dog with the MDR1 deletion mutation associated with ivermectin sensitivity. *J Am Vet Med Assoc* **223**: 1453-1455, 1434.
- Mellersh, C.S., C. Hitte, M. Richman, F. Vignaux, C. Priat, S. Jouquand, P. Werner, C. Andre, S. DeRose, D.F. Patterson et al. 2000. An integrated linkage-radiation hybrid map of the canine genome. *Mamm Genome* **11**: 120-130.
- Mosher, D.S., P. Quignon, C.D. Bustamante, N.B. Sutter, C.S. Mellersh, H.G. Parker, and E.A. Ostrander. 2007. A mutation in the myostatin gene increases muscle mass and enhances racing performance in heterozygote dogs. *PLoS Genet* **3**: e79.
- Neff, M.W. and J. Rine. 2006. A fetching model organism. *Cell* **124**: 229-231.
- Neff, M.W., K.R. Robertson, A.K. Wong, N. Safra, K.W. Broman, M. Slatkin, K.L. Mealey, and N.C. Pedersen. 2004. Breed distribution and history of canine *mdr1-1Delta*, a pharmacogenetic mutation that marks the emergence of breeds from the collie lineage. *Proc Natl Acad Sci U S A* **101**: 11725-11730.
- Nelson, O.L., E. Carsten, S.A. Bentjen, and K.L. Mealey. 2003. Ivermectin toxicity in an Australian Shepherd dog with the MDR1 mutation associated with ivermectin sensitivity in Collies. *J Vet Intern Med* **17**: 354-356.
- Ouahchi, K., N. Lindeman, and C. Lee. 2006. Copy number variants and pharmacogenomics. *Pharmacogenomics* **7**: 25-29.
- Parker, H.G., L.V. Kim, N.B. Sutter, S. Carlson, T.D. Lorentzen, T.B. Malek, G.S. Johnson, H.B. DeFrance, E.A. Ostrander, and L. Kruglyak. 2004. Genetic structure of the purebred domestic dog. *Science* **304**: 1160-1164.

- Parker, H.G., A.V. Kukekova, D.T. Akey, O. Goldstein, E.F. Kirkness, K.C. Baysac, D.S. Mosher, G.D. Aguirre, G.M. Acland, and E.A. Ostrander. 2007. Breed relationships facilitate fine-mapping studies: a 7.8-kb deletion cosegregates with Collie eye anomaly across multiple dog breeds. *Genome Res* **17**: 1562-1571.
- Patterson, D.F., M.E. Haskins, P.F. Jezyk, U. Giger, V.N. Meyers-Wallen, G. Aguirre, J.C. Fyfe, and J.H. Wolfe. 1988. Research on genetic diseases: reciprocal benefits to animals and man. *J Am Vet Med Assoc* **193**: 1131-1144.
- Perry, G.H., A. Ben-Dor, A. Tsalenko, N. Sampas, L. Rodriguez-Revenga, C.W. Tran, A. Scheffer, I. Steinfeld, P. Tsang, N.A. Yamada et al. 2008. The fine-scale and complex architecture of human copy-number variation. *Am J Hum Genet* **82**: 685-695.
- Perry, G.H., J. Tchinda, S.D. McGrath, J. Zhang, S.R. Picker, A.M. Caceres, A.J. Iafrate, C. Tyler-Smith, S.W. Scherer, E.E. Eichler et al. 2006. Hotspots for copy number variation in chimpanzees and humans. *Proc Natl Acad Sci U S A* **103**: 8006-8011.
- Pollinger, J.P., C.D. Bustamante, A. Fledel-Alon, S. Schmutz, M.M. Gray, and R.K. Wayne. 2005. Selective sweep mapping of genes with large phenotypic effects. *Genome Res* **15**: 1809-1819.
- Quignon, P., L. Herbin, E. Cadieu, E.F. Kirkness, B. Hedan, D.S. Mosher, F. Galibert, C. Andre, E.A. Ostrander, and C. Hitte. 2007. Canine population structure: assessment and impact of intra-breed stratification on SNP-based association studies. *PLoS ONE* **2**: e1324.
- Redon, R., S. Ishikawa, K.R. Fitch, L. Feuk, G.H. Perry, T.D. Andrews, H. Fiegler, M.H. Shapero, A.R. Carson, W. Chen et al. 2006. Global variation in copy number in the human genome. *Nature* **444**: 444-454.
- Rueda, O.M. and R. Diaz-Uriarte. 2007. Flexible and accurate detection of genomic copy-number changes from aCGH. *PLoS Comput Biol* **3**: e122.
- Salmon Hillbertz, N.H., M. Isaksson, E.K. Karlsson, E. Hellmen, G.R. Pielberg, P. Savolainen, C.M. Wade, H. von Euler, U. Gustafson, A. Hedhammar et al. 2007. Duplication of FGF3, FGF4, FGF19 and ORAOV1 causes hair ridge and predisposition to dermoid sinus in Ridgeback dogs. *Nat Genet* **39**: 1318-1320.
- Savolainen, P., Y.P. Zhang, J. Luo, J. Lundberg, and T. Leitner. 2002. Genetic evidence for an East Asian origin of domestic dogs. *Science* **298**: 1610-1613.
- Saxena, R., B.F. Voight, V. Lyssenko, N.P. Burt, P.I. de Bakker, H. Chen, J.J. Roix, S. Kathiresan, J.N. Hirschhorn, M.J. Daly et al. 2007. Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* **316**: 1331-1336.
- Schuetz, E.G. 2004. Lessons from the CYP3A4 promoter. *Mol Pharmacol* **65**: 279-281.
- Sebat, J. 2007. Major changes in our DNA lead to major changes in our thinking. *Nat Genet* **39**: S3-5.
- Sebat, J., B. Lakshmi, J. Troge, J. Alexander, J. Young, P. Lundin, S. Maner, H. Massa, M. Walker, M. Chi et al. 2004. Large-scale copy number polymorphism in the human genome. *Science* **305**: 525-528.
- Sharp, A.J., D.P. Locke, S.D. McGrath, Z. Cheng, J.A. Bailey, R.U. Vallente, L.M. Pertz, R.A. Clark, S. Schwartz, R. Graves et al. 2005. Segmental duplications and copy-number variation in the human genome. *Am J Hum Genet* **77**: 78-88.
- She, X., Z. Cheng, S. Zollner, D.M. Church, and E.E. Eichler. 2008. Mouse segmental duplication and copy number variation. *Nat Genet* **40**: 909-914.
- Sutter, N.B., C.D. Bustamante, K. Chase, M.M. Gray, K. Zhao, L. Zhu, B. Padhukasahasram, E. Karlins, S. Davis, P.G. Jones et al. 2007. A single IGF1 allele is a major determinant of small size in dogs. *Science* **316**: 112-115.
- Sutter, N.B. and E.A. Ostrander. 2004. Dog star rising: the canine genetic system. *Nat Rev Genet* **5**: 900-910.

- Trepanier, L.A. 2004. Idiosyncratic toxicity associated with potentiated sulfonamides in the dog. *J Vet Pharmacol Ther* **27**: 129-138.
- Tuzun, E., J.A. Bailey, and E.E. Eichler. 2004. Recent segmental duplications in the working draft assembly of the brown Norway rat. *Genome Res* **14**: 493-506.
- Tuzun, E., A.J. Sharp, J.A. Bailey, R. Kaul, V.A. Morrison, L.M. Pertz, E. Haugen, H. Hayden, D. Albertson, D. Pinkel et al. 2005. Fine-scale structural variation of the human genome. *Nat Genet* **37**: 727-732.
- Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. *Genome Biol* **4**: 231.
- Ventura, M., J.M. Mudge, V. Palumbo, S. Burn, E. Blennow, M. Pierluigi, R. Giorda, O. Zuffardi, N. Archidiacono, M.S. Jackson et al. 2003. Neocentromeres in 15q24-26 map to duplicons which flanked an ancestral centromere in 15q25. *Genome Res* **13**: 2059-2068.
- Vila, C., P. Savolainen, J.E. Maldonado, I.R. Amorim, J.E. Rice, R.L. Honeycutt, K.A. Crandall, J. Lundeberg, and R.K. Wayne. 1997. Multiple and ancient origins of the domestic dog. *Science* **276**: 1687-1689.
- Wagner, A.E., B.D. Wright, and P.W. Hellyer. 2003. Myths and misconceptions in small animal anesthesia. *J Am Vet Med Assoc* **223**: 1426-1432.
- Wayne, R.K. and E.A. Ostrander. 2007. Lessons learned from the dog genome. *Trends Genet* **23**: 557-567.

Table 1 Summary of CNVs identified in each sample.

| Breed | Number of CNVs | | | Gain | Loss | Average Size (kb) | Genes |
|-------------------------------|----------------|-----------|-----------|-----------|------------|-------------------|------------|
| | Total | ChrUn | Unique | | | | |
| Afghan Hound | 154 | 77 | 2 | 71 | 83 | 23.2 | 89 |
| Basenji | 298 | 106 | 22 | 103 | 195 | 34.7 | 216 |
| Belgian Shepherd Dog Tervuren | 185 | 41 | 10 | 109 | 76 | 36.5 | 130 |
| Doberman Pinscher | 220 | 61 | 2 | 102 | 118 | 38.8 | 138 |
| German Shepherd | 284 | 88 | 7 | 122 | 162 | 42.8 | 192 |
| German Shorthaired Pointer | 118 | 56 | 7 | 84 | 34 | 24.9 | 78 |
| Golden Retriever | 145 | 32 | 4 | 76 | 69 | 29.5 | 90 |
| Labrador Retriever | 249 | 112 | 27 | 97 | 152 | 24.9 | 131 |
| Pug | 131 | 91 | 43 | 75 | 56 | 31.8 | 38 |
| Rottweiler | 132 | 53 | 1 | 34 | 98 | 39.6 | 69 |
| Shetland Sheepdog | 259 | 76 | 10 | 70 | 189 | 36.3 | 182 |
| Siberian Husky | 277 | 95 | 16 | 128 | 149 | 34.2 | 171 |
| Standard Poodle | 219 | 72 | 21 | 106 | 113 | 37.1 | 147 |
| Boxer (Tasha) | 169 | 74 | 9 | 78 | 91 | 34.0 | 84 |
| West Highland White Terrier | 191 | 57 | 4 | 107 | 84 | 38.8 | 142 |
| Whippet | 125 | 84 | 6 | 86 | 39 | 24.7 | 68 |
| Wolf | 251 | 62 | 19 | 92 | 159 | 37.2 | 144 |
| Yorkshire Terrier | 175 | 45 | 7 | 38 | 137 | 39.6 | 121 |
| Self | 1 | 0 | 1 | 0 | 1 | 9.0 | 0 |
| Averages | 199 | 71 | 12 | 88 | 111 | 33.8 | 124 |

Figure Legends


- Figure 1** **The genomic architecture of canine segmental duplications and CNVs.**
Black lines represent all 38 canine autosomes, the X chromosome, and the uncharacterized chromosome (Un). Duplicated bases predicted by WGAC and WSSD are plotted as orange and blue rectangles below and above, respectively, each chromosome. Over 80% of chrUn contains duplicated bases. Of the autosomes, chromosomes 9, 16, and 26 possess the highest percentages of duplicated bases (over 4% of each chromosome) while chromosomes 12, 30, and 33 show the least amount of duplicated bases (less than 0.35% of each chromosome). Unique CNV regions (see text) are denoted by red rectangles.
- Figure 2** **Validation of duplicons by FISH analysis.** (A) Example of an interchromosomal duplication detected with clone CH82-381N09. (B) and (C) show two representative examples of tandem intrachromosomal duplication detected with clones CH82-381N09 and CH82-331L01, respectively.
- Figure 3** **Heatmap representation of CNVs.** Each row represents one of the 678 unique CNV regions and columns correspond to dogs. For each CNV region, boxes are colored as black, magenta, and green depending on whether the individual showed no copy number variation, a loss, or a gain, respectively. CNV regions that show both a loss and a gain within an individual dog (see text) are colored yellow. Horizontal white lines separate CNV calls from single copy control regions, and CNVs that exhibit only losses, gains, or both gains and losses. Within each class, CNV regions are sorted from low to high




frequency and from left to right dogs are sorted by decreasing number of CNVs. Dog breeds are abbreviated as follows: Basenji (BAS), Shetland Sheepdog (SHS), German Shepherd (GSH), Siberian Husky (SBH), Wolf (WLF), Labrador Retriever (LBR), Doberman Pinscher (DOB), Standard Poodle (STP), Belgian Shepherd Tervuren (BST), West Highland White Terrier (WST), Yorkshire Terrier (YRK), Boxer (Tasha) (TSH), Afghan Hound (AFH), Golden Retriever (GLR), Rottweiler (ROT), Pug (PUG), Whippet (WHP), German Short Haired Pointer (GSP), and Self-Self hybridization (SLF).

Figure 4 **qPCR of *GCKR* and *PHYH* regions.** Each plot shows the relative copy number in comparison to the reference (y-axis) for each breed (x-axis). The reference sample was the same Boxer that was used as the reference in the aCGH experiments. Note, because the *GCKR* and *PHYH* regions are located in segmental duplications a gain or loss is not expected to yield a relative copy change of 2 and 0.5, respectively. For example, if the reference sample contains three copies, a gain in the test sample would result in an expected relative copy number of 1.33. Vertical bars delimit 95% confidence intervals based on six independent replicates. False positives (CNVs predicted in the aCGH data but not confirmed by qPCR) are colored in red. Breed abbreviations are described in Figure 3.

Figure 5 **Example of a complex CNV region.** Distribution of \log_2 probe intensities across a 400 kb region of chromosome 17 that shows substantial variation in breakpoints across individuals and spatial heterogeneity of copy number

within individuals. Black, green, and magenta indicate regions called as no copy number variation, gains, and losses, respectively. The gene structure of *GCKR* is shown, with black lines and red boxes corresponding to introns and exons, respectively. For clarity, additional RefSeq genes are not shown. Regions with significant WGAC or WSSD scores are indicated by gray rectangles. Breeds are abbreviated as described in Figure 3.





CNVs

